



UNIFORM
CONVERGENCE
OF SEQUENCE &
SERIES
2019-20

An Extramural Seminar by
Dr Arun Kumar Tripathy,
Lecturer in Mathematics,
SSB College, Mahakalpara

**Organized by Department of
Mathematics, Pattamundai
College, Pattamundai**

On 22.02.2020

REPORT

A seminar was organised by Department of Mathematics, Pattamundai College, Pattamundai on 22.02.2020 on the topic "**UNIFORM CONVERGENCE OF SEQUENCE & SERIES**". Dr Arun Kumar Tripathy , Lecturer in Mathematics, S.S.B College, Mahakalapada who graced the seminar with his analytical thinking. We were able to get the beautiful glimpses of the students of our Department. Sri Arabinda Pandab, Head of the Department gave a key note address of the topic and welcomed the guests on the diace and the participants. The meeting was ended with a vote of thanks by Dr Nirmal Kumar Sahoo, another faculty member.

Uniform Convergence of Sequences and Series

-Dr Arun Kumar Tripathy, SSB College, Mahakalpara

1. UNIFORM CONVERGENCE OF SEQUENCES OF FUNCTIONS

Definition (Pointwise convergence). A sequence $\{f_n\}$ of functions *converges pointwise on A* to a function f if for any given $x \in A$ and $\varepsilon > 0$ there exists a $N = N(x, \varepsilon) \in \mathbb{N}$ such that

$$n > N \implies |f_n(x) - f(x)| < \varepsilon.$$

We saw in class that pointwise limits do not interact well with the concepts of continuity, integrability and differentiability from calculus. For this reason we introduce the following *stronger* notion of convergence.

Definition (Uniform convergence). A sequence $\{f_n\}$ of functions *converges uniformly on A* to a function f if for any given $\varepsilon > 0$ there exists a $N = N(\varepsilon) \in \mathbb{N}$ such that

$$n > N \implies |f_n(x) - f(x)| < \varepsilon, \text{ for every } x \in A.$$

The following Theorems illustrate that in contrast to pointwise convergence, *uniform convergence* interacts well with the concepts of calculus.

Theorem 1.1 (Uniform convergence preserves continuity). *If $f_n \rightarrow f$ uniformly on A and each function f_n is continuous at a point $a \in A$, then the limit function f is also continuous at a.*

Note: Symbolically we can express this result as

$$\lim_{x \rightarrow a} \lim_{n \rightarrow \infty} f_n(x) = \lim_{n \rightarrow \infty} \lim_{x \rightarrow a} f_n(x).$$

Theorem 1.2 (Integration preserves uniform convergence). *Suppose that $f_n \rightarrow f$ uniformly on $A = [a, b]$ and each function f_n is continuous on A. If for a fixed $x_0 \in A$ we define*

$$F_n(x) = \int_{x_0}^x f_n(t) dt \quad \text{and} \quad F(x) = \int_{x_0}^x f(t) dt$$

for all $x \in A$, then $F_n \rightarrow F$ uniformly on A.

Note: Symbolically we can express this result as

$$\lim_{n \rightarrow \infty} \int_{x_0}^x f_n(t) dt = \int_{x_0}^x \lim_{n \rightarrow \infty} f_n(t) dt.$$

Note that it is only as a consequence of Theorem 1.1 that the definition of the function F above makes sense.

Unfortunately, differentiation does not preserve uniform convergence; recall that the sequence of functions $f_n(x) = \frac{1}{n} \sin(n^2 x)$ converges to 0 uniformly, but the derived sequence $\{f'_n\}$ does not even converge pointwise! The following result follows from Theorem 1.2 (and the Fundamental Theorem of Calculus).

Theorem 1.3 (Differentiation). *Let $\{f_n\}$ be a sequence of functions on $A = [a, b]$ such that each f_n is differentiable on A with continuous derivative. If $\lim_{n \rightarrow \infty} f_n(x_0)$ exists for some $x_0 \in A$ and f'_n converges uniformly on A to some function g , then f_n converges uniformly on A to a differentiable function f with $f' = g$.*

Note: Symbolically we can express this result as

$$\frac{d}{dx} \lim_{n \rightarrow \infty} f_n(x) = \lim_{n \rightarrow \infty} \frac{d}{dx} f_n(x).$$

2. UNIFORM CONVERGENCE OF INFINITE SERIES OF FUNCTIONS

We can apply this notion of convergence (and hence the above results) to series of functions.

Definition (Uniform convergence of series). A series $\sum_{n=0}^{\infty} f_n$ of functions converges uniformly on A to a function s if the sequence of partial sums $\{s_n\}$ of the series converges uniformly on A to s .

The following result gives a sufficient condition for uniform convergence.

Weierstrass M-test. If $|f_n(x)| \leq M_n$ for each $x \in A$ and the sequence $\{M_n\}$ is summable, then the series $\sum_{n=0}^{\infty} f_n$ converges uniformly on A .

The following results follow by applying Theorems 1.1, 1.2 and 1.3 respectively to the sequence of partial sums $s_n(x) = \sum_{k=0}^n f_k(x)$.

Theorem 2.1. If $\sum_{n=0}^{\infty} f_n$ converges uniformly on A and each term f_n is continuous at a point $a \in A$, then the sum function s is also continuous at a .

Note: Symbolically we can express this result as

$$\lim_{x \rightarrow a} \sum_{n=0}^{\infty} f_n(x) = \sum_{n=0}^{\infty} \lim_{x \rightarrow a} f_n(x).$$

Theorem 2.2. Suppose that $\sum_{n=0}^{\infty} f_n$ converges uniformly on $A = [a, b]$ to a sum function s where each term f_n is continuous on A . If for a fixed $x_0 \in A$ we define

$$S_n(x) = \int_{x_0}^x s_n(t) dt \left(= \sum_{k=0}^n \int_{x_0}^x f_k(t) dt \right) \quad \text{and} \quad S(x) = \int_{x_0}^x s(t) dt$$

for all $x \in A$, then $S_n \rightarrow S$ uniformly on A .

Note: Symbolically we can express this result as

$$\sum_{n=0}^{\infty} \int_{x_0}^x f_n(t) dt = \int_{x_0}^x \sum_{n=0}^{\infty} f_n(t) dt.$$

Again it is only as a consequence of Theorem 2.1 that the definition of the function S above makes sense.

Unfortunately, "term-by-term" differentiation may destroy convergence, even if the original series converges uniformly; the series $\sum_{n=0}^{\infty} (\sin nx)/n^2$ converges uniformly for all x by the M-test, but the series obtained by differentiating "term-by-term" is $\sum_{n=0}^{\infty} (\cos nx)/n$, which converges if and only if x is not an integer multiple of 2π .

Theorem 2.3. Let $\{f_n\}$ be a sequence of functions on $A = [a, b]$ such that each f_n is differentiable on A with continuous derivative. If $\sum_{n=0}^{\infty} f_n(x_0)$ converges for some $x_0 \in A$ and the series of derivatives $\sum_{n=0}^{\infty} f'_n$ converges uniformly on A to a sum function g , then $\sum_{n=0}^{\infty} f_n$ converges uniformly on A to a differentiable sum function s with $s' = g$.

Note: Symbolically we can express this result as

$$\frac{d}{dx} \sum_{n=0}^{\infty} f_n(x) = \sum_{n=0}^{\infty} \frac{d}{dx} f_n(x).$$

Note that the problem of interchanging the operations of differentiation and summation is, in general, more serious than in the case of integration. Manipulations that we are familiar with carrying out on finite sums do not always carry over to infinite series, even if the series involved are uniformly convergent. We turn next to special series of functions, known as power series, which can be manipulated in many respects as though they were finite sums.

3. POWER SERIES

Definition. A power series (centered at a) is a series of the form $\sum_{n=0}^{\infty} a_n(x-a)^n$ where a is some fixed real number, its domain of convergence is given by $\{x \in \mathbb{R} \mid \sum_{n=0}^{\infty} a_n(x-a)^n \text{ converges}\}$.

Proposition 3.1. If the power series $\sum_{n=0}^{\infty} a_n(x-a)^n$ converges for $x-a=c$, then it converges absolutely for all x such that $|x-a| < |c|$.

Corollary 3.2. The domain of convergence of any power series centered at a is an interval centered at a , it may, however, include neither, one, or both of the endpoints of the interval.

Definition. If R is a nonnegative real number such that the power series $\sum_{n=0}^{\infty} a_n(x-a)^n$ converges for all real x with $|x-a| < R$ and diverges for all real x with $|x-a| > R$ then R is called the radius of convergence for the power series. If the power series converges for all real x , then we say that the radius of convergence is infinite.

Proposition 3.3. If $\lim_{n \rightarrow \infty} |a_n/a_{n+1}| = R$ exists as a finite number or is infinite, then R is the radius of convergence for the power series $\sum_{n=0}^{\infty} a_n(x-a)^n$.

The following proposition is an immediate consequence of the Weierstrass M-test.

Proposition 3.4. If the power series $\sum_{n=0}^{\infty} a_n(x-a)^n$ converges for all $|x-a| < R$, then for any $0 < c < R$ it converges uniformly on the interval $[a-c, a+c]$.

In summary, it follows from Proposition 3.1 that if the power series $\sum_{n=0}^{\infty} a_n(x-a)^n$ has a radius of convergence R , then it converges absolutely on the open interval $(a-R, a+R)$, and from Proposition 3.4 that it converges uniformly on every closed interval $[a-c, a+c]$, where $0 < c < R$.

Proposition 3.5 (Properties of power series). Let $\sum_{n=0}^{\infty} a_n(x-a)^n$ have radius of convergence $R > 0$ and suppose $\sum_{n=0}^{\infty} b_n(x-a)^n$ also converges for $x \in A = (a-R, a+R)$. Define $f: A \rightarrow \mathbb{R}$ and $g: A \rightarrow \mathbb{R}$ by

$$f(x) = \sum_{n=0}^{\infty} a_n(x-a)^n \quad \text{and} \quad g(x) = \sum_{n=0}^{\infty} b_n(x-a)^n.$$

1. (Arithmetic) If $x \in A$, then

$$(a) \quad \lambda f(x) = \sum_{n=0}^{\infty} \lambda a_n(x-a)^n \quad \text{for all } \lambda \in \mathbb{R}$$

$$(b) \quad f(x) + g(x) = \sum_{n=0}^{\infty} (a_n + b_n)(x-a)^n$$

$$(c) \quad f(x)g(x) = \sum_{n=0}^{\infty} \left(\sum_{k=0}^n a_k b_{n-k} \right) (x-a)^n$$

2. (Calculus) On $A = (a-R, a+R)$, f is differentiable (and hence continuous) and, for $x \in A$,

$$f'(x) = \sum_{n=0}^{\infty} n a_n (x-a)^{n-1} \quad \text{and} \quad \int_a^x f(t) dt = \sum_{n=0}^{\infty} a_n \frac{(x-a)^{n+1}}{n+1}.$$

In particular these series also have radius of convergence R .

3. (Equating Coefficients) If $f = g$, then $a_n = b_n = \frac{f^{(n)}(a)}{n!}$ for all $n \in \mathbb{N}$.

The second set of results above follow from the corresponding results in section 2, recall that we actually proved the differentiation result above a little more directly.

4. TAYLOR SERIES

Applying Proposition 3.5 (part 2) to the geometric series we are able to establish the validity of the expressions

$$\ln(1+x) = \sum_{n=1}^{\infty} \frac{(-1)^{n+1} x^n}{n} \quad \text{and} \quad \tan^{-1} x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{2n+1}$$

for all $|x| < 1$. The above series actually converge for $x = 1$ and $|x| = 1$ respectively, do they also converge to the functions there?

Another consequence of Proposition 3.5 (part 2) is that the sum function of a power series has derivatives of every order, these are obtained by repeated term by term differentiation of the power series. Hence if a given function f has a power series expansion then it must take the form

$$\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x-a)^n.$$

This is called the *Taylor series generated by f at a* . The n th partial sum

$$P_n^f(x) = \sum_{k=0}^n \frac{f^{(k)}(a)}{k!} (x-a)^k$$

is called the *n th order Taylor polynomial of f at a* .

Two questions naturally arise: Given a function f , does its Taylor series converge for any x other than $x = a$? If so, is its sum equal to $f(x)$? The answer to both these questions is, in general, no.

Example. The Maclaurin series of the function

$$f(x) = \begin{cases} e^{-1/x^2} & x \neq 0 \\ 0 & x = 0 \end{cases}$$

is identically 0, and hence does not converge to f for any $x \neq 0$.

Necessary and sufficient conditions for answering both questions in the affirmative can be expressed in terms of

$$R_n^f(x) := f(x) - P_n^f(x),$$

this is called the *n th remainder of f about a* .

Proposition 4.1. $f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x-a)^n \iff P_n^f(x) \rightarrow f(x) \iff R_n^f(x) \rightarrow 0$.

Theorem 4.2 (Taylor's Theorem). *Suppose that $f, f', \dots, f^{(n)}$ are continuous on $[\alpha, \beta]$ and $f^{(n)}$ is differentiable on (α, β) , then for any x and a in $[\alpha, \beta]$ there exists a number c between x and a so that*

$$R_n^f(x) = \frac{f^{(n+1)}(c)}{(n+1)!} (x-a)^{n+1}.$$

We finish this section with a result which answers in the affirmative the first question we posed.

Theorem 4.3 (Abel's Theorem). *Suppose that $\sum_{n=0}^{\infty} a_n$ converges and $f(x) = \sum_{n=0}^{\infty} a_n x^n$, for $|x| < 1$, then $\lim_{x \rightarrow 1^-} f(x) = \sum_{n=0}^{\infty} a_n$.*

Standard Maclaurin Series (Taylor series centered at 0)

1. $(1-x)^{-1} = \sum_{n=0}^{\infty} x^n \quad (-1 < x < 1)$
2. $e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} \quad (x \in \mathbb{R})$
3. $\ln(1+x) = \sum_{n=1}^{\infty} \frac{(-1)^{n+1} x^n}{n} \quad (-1 < x \leq 1)$
4. $\tan^{-1} x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{2n+1} \quad (-1 \leq x \leq 1)$
5. $\cos x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!} \quad (x \in \mathbb{R})$
6. $\sin x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!} \quad (x \in \mathbb{R})$

7. Binomial Series: For $r \in \mathbb{R}$

$$(1+x)^r = \sum_{n=0}^{\infty} \binom{r}{n} x^n \quad (x \in A)$$

where $\binom{r}{0} = 1$ and, for $n \in \mathbb{N}$,

$$\binom{r}{n} = \frac{r(r-1)\cdots(r-n+1)}{n!}.$$

The set A depends on r . When $r \in \mathbb{N}$, then $A = \mathbb{R}$. In all cases, $(-1, 1) \subseteq A$.

5. OPERATIONS ON TAYLOR POLYNOMIALS AND SERIES (INFORMAL)

Throughout this section f is assumed to have derivatives of all orders on the open interval $(-R, R)$.

We recall that by definition, the n th order Maclaurin polynomial of f is given by

$$P_n^f(x) = \sum_{k=0}^n \frac{f^{(k)}(0)}{k!} x^k$$

and is related to the n th Maclaurin remainder of f by

$$f(x) - P_n^f(x) = R_n^f(x).$$

The following results are useful when performing calculations involving Taylor series. The first result allows us to reduce our discussions to simply Maclaurin series.

Proposition 5.1 (Translation). *Let $a \in \mathbb{R}$, and define a new function g by $g(x) = f(x-a)$. Then the n th order Taylor polynomial of g about a is $P_n^f(x-a)$ and n th order remainder of g is $R_n^f(x-a)$.*

The following results concern the calculus of Maclaurin polynomials.

Proposition 5.2 (Differentiation). *The n th order Maclaurin polynomial of f' is the derivative of $P_{n+1}^f(x)$ and n th order remainder of f' is the derivative of $R_{n+1}^f(x)$.*

Proposition 5.3 (Integration). *If $F(x) = \int_0^x f(t) dt$, for $|x| < R$, then the n th order Maclaurin polynomial of F is $P_n^F(x) = \int_0^x P_{n-1}^f(t) dt$ and n th order remainder of F is $R_n^F(x) = \int_0^x R_{n-1}^f(t) dt$.*

The result below is useful if you have to find Maclaurin polynomials or remainders of functions like $g(x) = x \sin(x^2)$, for example.

Proposition 5.4 (Monomial multiplication and composition). *Suppose that k and ℓ are positive integers and λ is a real number. If $g(x) = x^k f(\lambda x^\ell)$, then the $(n\ell+k)$ th order Maclaurin polynomial of g is $P_{n\ell+k}^g(x) = x^k P_n^f(\lambda x^\ell)$ and $(n\ell+k)$ th order remainder of g is $R_{n\ell+k}^g(x) = x^k R_n^f(\lambda x^\ell)$.*



OFFICE OF THE PRINCIPAL

Mobile : 9437376724

PATTAMUNDAI COLLEGE

NAAC ACCREDITED B+ GRADE

PATTAMUNDAI, KENDRAPARA, ODISHA - 754215

Ref No. : 318

Date..... 19/02/2020

To

Dr Arun Kumar Tripathy,
Lecturer in Mathematics,
S.S.B College, Mahakalpara.

Sub: - An invitation as Resource Person in the Extramural Seminar in
Department of Mathematics.

Sir,

It is my pleasure to invite you as **Resource Person** in the Extramural Seminar on the topic "**Uniform Convergence of Sequence & Series**" to be organised by Department of Mathematics, at 11.00 am on 22nd February 2020 in our institution.

Your kind presence for this occasion is highly solicited.

Yours Faithfully,


19.2.2020

Principal
Pattamundai College,

Principal
Pattamundai College

An Extramural Seminar on "Uniform Convergence of Sequence & Series"

Department of Mathematics

Pattamundai College, Pattamundai

Dt. 22.02.2020

Sl No	Name of the Resource Person/Faculty Member	Signature & Mobile Number
1	Dr Arun Kumar Tripathy	Arun Kumar Tripathy 9937596346
2	Arabinda Panda	Arabinda Panda 22/2/20 9937251024
3	Dr Nirmalakumar Sahoo	Nirmala Kumar Sahoo 8974405754
4		
5		
6		
7		
8		
9		
10		

An Extramural Seminar on "Uniform Convergence of Sequence & Series"

Department of Mathematics

Pattamundai College, Pattamundai

Dt. 22.02.2020

Sl No	Name of the Student	Roll Number	Signature & Mobile Number
1	Hihir Ranjan Tripathy	BSCP) 19-059	Hihir Ranjan Tripathy 9937890859
2	Sudhin Kanta Das	BS(P) 19-005	Sudhin Kanta Das 6370180517
3	Chandan senapati	BS (P) 19-118	Chandan senapati 6372898627
4	Manoj Kanta Sahoo	BS(P) 19-060	Manoj Kantasahoo 9861219798
5	Manasi Mohanty	BS-17-119	Manasi Mohanty 977076894
6	Rajanikanta Jena	BS-17-069	Rajanikanta Jena 8328920601
7	Debabrata Mahata	BS-17-158	Debabrata Mahata 9668795438
8	Kabita Patra	BS(P)-19-074	Kabita Patra 9861005124
9	Jyotiprava Sahoo	BS(P) 19-036	Jyotiprava Sahoo 8917469648
10	Aryabartini Jena	BS(P) 19-053	Aryabartini Jena 9668289747
11	Swagatika Nayak	BS17-019	Swagatika 7377738952.
12	Swara; KemaR Das	BS17-068	Swara 7328021411

13	Ansuman Sahoo	BS14-125	Ansuman	#325991657
14	Sushakanta Rauty	BSCP19-026	Sushakanta Rauty	
15	Depak Kumar Pani	BS(P)19-056	Depak Kumar Pani	
16	Goutam Behera	BSCP19-065	Goutam Behera	
17	Sai Prasad Swain	BS(P)19-100	Sai Prasad Swain	
18	Shruti Keshari Satapathy	BS(P)19-114	Shruti Keshari Satapathy	
19	Nandakishore Rout	BS-17-039	Nandakishore Rout	
20	Anuradha Rout	BS-18-074	Anuradha Rout	
21	Digantika Das	BS-18-140	Digantika Das	
22	Sheela Mahanta	BS18-064	Sheela Mahanta	
23	Saswotika Rath	BS-18-128	Saswotika Rath	
24	Manisha Guain	BS-18-139	Manisha Guain	
25	Elina Sween	BS-18-138	Elina Sween	
26	Usharani Mohanty	BS-18-080	Usharani Mohanty	
27	Angasha Pasola	BS18-004	Angasha Pasola	
28	Ashrita Bhuyan	BS18-026	Ashrita Bhuyan	
29	Kaivalash chandrea Malik	BS18-075	Kaivalash chandrea Malik	8847893009

30	Yamunpate Roy	BS18-105	Yamunpate Roy	637365579
31	Malaya Ranjan Swamy	BS18-070	Malaya Ranjan Swamy	6370492519
32	Raghavendra Biswal	BS18-045	Raghavendra Biswal	6271980094
33	Satya Narayan Das	BS-18-091	Satya Narayan Das	7077991009
34	Subrajit Rout	BS-18-065	Subrajit Rout	8144528007
35	Suchansu sekhar Rout	BS-17-111	Suchansu sekhar Rout	7978827314
36	Sreastimajani pradhan	BS17-065	Sreastimajani pradhan	6977764578
37	Rajain Barik	BS17-145	Rajain Barik	7853014013
38				
39				
40				
41				
42				
43				
44				
45				
46				

